Neuroscience Letters, 77 (1987) 367–371
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Adham, N. and Smith, M.E, Exposure of receptor sites on rat muscle membranes by a neurally released factor, 283
Agnati, L.F., see Citra, A., 25
Akasu, T., see Tsurusaki, M., 303
Akiyama, H., see Kaneko, T., 171
Amenta, D., Cavallotti, C. and Amenta, F., Dopamine receptors mediating the stimulation and the inhibition of adenylate cyclase in rat prostate gland, 66
Amenta, F., see Amenta, D., 66
Artalejo, C.R., see Gandía, L., 333
Asin. K.E., see Wirtshafter, D., 261

Baudyšová, see Bolotina, V., 298
Benistant, C., Rey, C., Fonlupt, P. and Pacheco, H., Increase of in vitro brain [^{3}H]muscimol binding after L-homocysteine administration to rat, 237
Benmansour, S., Bonnet, J.-J., Protais, P. and Costentin, J., Sodium independence of the binding of [^{3}H]GBR 12783 and other dopamine uptake inhibitors to the dopamine uptake complex, 97
Bérod, A., see Kitahama, K., 155
Beuter, A., Carrière, L. and Boucher, J.P., Relationships between electromyography and kinematics in human stepping strategies, 119
Binet, S., Cohen, E. and Meininger, V., Heterogeneity of cold-stable and cold-labile tubulin in axon- and soma-enriched portions of the adult mous brain, 166
Bissette, G., see Widerlöv, E., 311
Blakemore, W.F., Crang, A.J., Evans, R.J. and Patterson, R.C., Rat Schwann cell remyelination of demyelinated cat CNS axons: evidence that injection of cell suspensions of CNS tissue results in Schwann cell remyelination, 15
Blakemore, W.F. Crang, A.J. and Patterson, R.C., Schwann cell remyelination of CNS axons following injection of cultures of CNS cells into areas of persistant demyelination, 20
Bolioli, B., see Dajas, F., 87
Bolotina, V., Borecký, J., Vlachová, V., Baudyšová, M. and Vyskočil, F., Voltage-dependent chloride channels with several substates in excised patches from mouse neuroblastoma cells, 298
Bonnet, J.-J., see Benmansour, S., 97
Borecký, J., see Bolotina, V., 298
Boucher, J.P., see Beuter, A., 119
Bouvet, J.F., Delaleu, J.C. and Holley, A., Olfactory receptor cell function is affected by trigeminal nerve activity, 181
Brown, D.A., see Marrion, N.V., 55
Buu, N.T., see Debinski, W., 92

Calas, A., see Decavel, C., 149
Campistron, G., see Esclapez, M., 131
Cantin, M., see Debinski, W., 92
Carlsson, K.H. and Jurna, I., Central analgesic effect of paracetamol manifested by depression of nociceptive activity in thalamic neurones of the rat, 339
Carobi, C., Capsaicin-sensitive vagal afferent neurons innervating the rat pancreas, 5
Carrière, L., see Beuter, A., 119
Castello, M.E., see Dajas, F., 87
Cavallotti, C., see Amenta, 66
Chalmers, J., see Hodgson, A., 125
Chrachri, A. and Clarac, F., Induction of rhytmic activity in motoneurons of crayfish thoracic ganglia by cholinergic agonists, 49
Cicero, S., see Facchinetti, F., 349
Cintra, A., Fuxe, K., Härfstrand, A., Agnati, L.F., Wikström, A.-C., Okret, S., Vale, W. and Gustafsson, J.-Å., Presence of glucocorticoid receptor immunoreactivity in corticotrophin releasing factor and in growth hormone releasing factor immunoreactive neurons of the rat di- and telencephalon, 25
Clarac, F., see Chrachri, A., 49
Cohen, E., see Binet, S., 166
Colebatch, J.G. and Porter, R., 'Long-latency' responses occurring with stratle in the conscious monkey, 43
Costentin, J., see Benmansour, S., 97
Couraud, J.-Y., see Pollard, H., 267
Crang, A.J., see Blakemore, W.F., 15
Crang, A.J., see Blakemore, W.F., 20
Curti, M., see Fussi, F., 308
Czachurski, J., see Dembowsky, K., 31

Dajas, F., Bolioli, B., Castello, M.E. and Silveira, R., Rat striatal acetylcholinesterase inhibition by Fasciculin (a polypeptide from green mamba snake venom), 87
Dantzer, R., see Le Moal, M., 353
Debinski, W., Kuchel, O., Buu, N.T., Cantin, M. and Genest, J., Atrial natriuretic factor partially inhibits the stimulated catecholamine synthesis in superior carvical ganglia of the rat, 92
Decavel, C., Geffard, M. and Calas, A., Comparative study of dopamine- and noradrenaline-immunoreactive terminals in the paraventricular and supraoptic nuclei of the rat, 149
De Jong, W., see Van den Buuse, M., 71
DelMastro, R., see Sabel, B.A., 360
Delaleu, J.C., see Bouvet, J.F., 181
Dembowsky, K., Czachurski, J. and Seller, H., Rostrocaudal location of sympathetic preganglionic neurones within the third thoracic segment of the cat spinal cord investigated by the retrograde transport of horseradish peroxidase and by recording of antidromic field potentials, 31
Denoyer, M., see Kitahama, K., 155
De Witte, P.A., see Laduron, P.M., 344
Dietl, H., Regulation of the release of neuronally localized corticosterone in rat forebrain areas by catecholamines, 76
Dudel, J., see Franke, Ch., 199
Dun, N.J., see Mo, N., 327
Dunbar, G.L., see Sabel, B.A., 360

Esclapez, M., Campistron, G. and Trottler, S., Immunocytochemical localization and morphology of GABA-containing neurons in the prefrontal and frontoparietal cortex of the rat, 131
Evans, R.J., see Blakemore, W.F., 15

Facchinetti, F., Petraglia, F., Cicero, S., Nappi, G., Consoli, V. and Genazzani, A.R., No gradient exists between lumbar and ventricular cerebrospinal fluid β-endorphin, 349
Falk, G., see Shiells, R.A., 221
Fonlupt, P., see Benistant, C., 237
Fonteríz, R.I., see Gandía, L., 333
Franke, Ch., Hatt, H. and Dudel, J., Liquid filament switch for ultra-fast exchanges of solutions at excised patches of synaptic membrane of crayfish muscle, 199
Fritzsch, B., see Sonntag, R., 143
Fussi, F., Savoldi, F. and Curti, M., Identification of *N*-carboxyethyl γ-aminobutyric acid in bovine brain and human cerebrospinal fluid, 308
Fuxe, K., see Cintra, A., 25

Gandía, L., López, M.G., Fonteríz, R.I., Artalejo, C.R. and García, A.G., Relative sensitivities of chromaffin cell calcium channels to organic and inorganic calcium antagonists, 333
García, A.G., see Gandía, L., 333
Geffard, M., see Decavel, C., 149
Genazzani, A.R., see Facchinetti, F., 349
Genest, J., see Debinski, W., 92
Gustafsson, J.-Å., see Cintra, A., 25

Hamashima, Y., see Yoshioka, H., 10
Härfstrand, A., see Cintra, A., 25
Hatt, H., see Franke, Ch., 199
Heizmann, C.W., see Röhrenbeck, J., 255
Hess, G., Kuhnt, U. and Voronin, L.L., Quantal analysis of paired-pulse facilitation in guinea pig hippocampal slices, 187
Hodgson, A., Minson, J., Chalmers, J. and Pilowsky, P., The use of microinjected colloidal gold and immunocytochemistry to localise pressor sites in the rostral medulla oblongata of the rat, 125
Holley, A., see Bouvet, J.F., 181

Ishiki, R., see Naoi, M., 215
Ito, K.-i., see Izumi, Y., 176
Iyengar, S., Kim, H.S. and Wood, P.L., Agonist action of the agonist/antagonist analgesic butorphanol on dopamine metabolism in the nucleus accumbens of the rat, 226
Izumi, Y., Ito, K.-i., Miyakawa, H. and Kato, H., Requirement of extracellular Ca^{2+} after tetanus for inducation of long-term potentiation in guinea pig hippocampal slices, 176

Janssen, P.F.M., see Scherman, D., 231
Johnston, G.A.R., see Ong, J., 109
Jordan, F.L. and Thomas, W.E., Identification of somatostatin-containing neurons in primary cultures of rat cerebral cortex, 249
Jouvet, M., see Kitahama, K., 155
Jurna, I., see Carlsson, K.H., 339

Kamiya, H., see Nagashima, A., 103
Kaneko, T., Akiyama, H. and Mizuno, N., Immunohistochemical demonstration of glutamate dehydrogenase in astrocytes, 171
Kato, H., see Izumi, Y., 176
Kayama, Y. and Ogawa, T., Electrophysiology of ascending, possibly cholinergic neurons in the rat laterodorsal tegmental nucleus: comparison

with monoamine neurons, 277
Kerr, D.I.B., see Ong, J., 109
Kim, H.S., see Iyengar, S., 226
Kimura, T., see Yoshioka, H., 10
Kitahama, K., Bérod, A., Denoyer, M. and Jouvet, M., Visualization of tyrosine hyroxylase-immunoreactive neurons in the cat dorsal motor vagal cells after treatment with parachlorophenylalanine, 155
Koob, G.F., see Le Moal, M., 353
Krnjević, K., see Zhang, L., 287
Krukoff, T.L., NPY-containing nerve terminals and fibers in close association with peptidergic sympathetic preganglionic neurons, 1
Kuchel, O., see Debinski, W., 92
Kuhnt, U., see Hess, G., 187
Kurosawa, M., Sato, A., Sato, Y. and Suzuki, H., Undiminished reflex responses of adrenal sympathetic nerve activity to stimulation of baroreceptors and cutaneous mechanoreceptors in aged rats, 193

Laduron, P.M. and De Witte, P.A., Enhanced axonal transport of receptor-bound opiate in ethanol-treated rats, 344
Laduron, P.M., see Scherman, D., 231
Lee, A.Y.S, see Wong, T.M., 61
Le Moal, M., Dantzer, R., Michaud, B. and Koob, G.F., Centrally injected arginine vasopressin (AVP) facilitates social memory in rats, 353
Lemon, R.N., see Mantel, G.W.H., 113
Licata, F., Li Volsi, G., Maugeri, G. and Santangelo, F., Integration of cortical and peripheral information in the lateral vestibular nucleus in the cat, 293
Livett, B.G., see Marley, P.D., 81
Li Volsi, G., see Licata, F., 293
Llorens-Cortes, C., see Pollard, H., 267
Lopes da Silva, F.H., see Melchers, B.P.C., 37
López, M.G., see Gandía, L., 333
Lukowiak, K., A blood-borne factor from food-satiated *Aplysia*, suppresses the gill withdrawal reflex in in vitro preparations from unsatiated animals, 205

Mantel, G.W.H. and Lemon, R.N., Cross-correlation reveals facilitation of single motor units in thenar muscles by single corticospinal neurones in the conscious monkey, 113
Marley, P.D. and Livett, B.G., Differences between the mechanisms of adrenaline and noradrenaline secretion from isolated, bovine, adrenal chromaffin cells, 81
Marrion, N.V., Smart, T.G. and Brown, D.A., Membrane currents in adult rat superior cervical ganglia in dissociated tissue culture, 55
Masui, H., see Nagashima, A., 103
Matesz, C., see Székely, G., 161
Maugeri, G., see Licata, F., 293
Meininger, V., see Binet, S., 166
Melchers, B.P.C., Pennartz, C.M.A. and Lopes da Silva, F.H., Differential effects of elevated extracellular calcium concentrations on field potentials in dentate gyrus and CA_1 of the rat hippocampal slice preparation, 37
Michaud, B., see Le Moal, M., 353
Minson, J., see Hodgson, A., 125
Miyakawa, H., see Izumi, Y., 176
Miyake, T., see Yoshioka, H., 10
Mizuno, N. see Kaneko, T., 171
Mo, N. and Dun, N.J., Excitatory postsynaptic potentials in neonatal rat sympathetic preganglionic neurons: possible mediation by NMDA receptors, 327
Müller, C.M., γ-Aminobutyric acid immunoreactivity in brainstem auditory nuclei of the chicken, 272

Nagashima, A., Takano, Y., Masui, H. and Kamiya, H., Evidence that neurokinin A (substance K) neurons project from the striatum to the substantia nigra in rats, 103
Nagatsu, T., see Naoi, M., 215
Naoi, M., Nomura, Y., Ishiki, R., Suzuki, H. and Nagatsu, T., 1,4-Benzoquinone as a new inhibitor of monoamine oxidase, 215
Nappi, G., see Facchinetti, F., 349
Nemeroff, C.B., see Widerlöv, E., 311
Nguyen, P., see Pomeranz, B., 316
Nishino, K., see Yoshioka, H., 10
Nomura, Y., see Naoi, M., 215

Ogawa, T., see Kayama, Y., 277
Ohshio, G., see Yoshioka, H., 10
Okret, S., see Cintra, A., 25
Ong, J., Kerr, D.I.B. and Johnston, G.A.R., Differing actions of β-phenyl-GABA and baclofen in the guinea pig isolated ileum, 109

Pacheco, H., see Benistant, C., 237
Patterson, R.C., see Blakemore, W.F., 15

Patterson, R.C., see Blakemore, W.F., 20
Pennartz, C.M.A., see Melchers, B.P.C., 37
Petraglia, F., see Facchinetti, F., 349
Pilowsky, P., see Hodgson, A., 125
Pollard, H., Llorens-Cortes, C., Couraud, J.-Y., Ronco, P., Verroust, P. and Schwartz, J.-C., Enkephalinase (EC 3.4.24.11) is highly localized to a striatonigral pathway in rat brain, 267
Pomeranz, B. and Nguyen, P., Intrathecal diazepam suppresses nociceptive reflexes and potentiates electroacupuncture effects in pentobarbital-anesthetized rats, 316
Porter, R., see Colebatch, J.G., 43
Potter, E.K., Cardiac vagal action and plasma levels of neuropeptide Y following intravenous injection in the dog, 243
Protais, P., see Benmansour, S., 97

Reiner, P.B., Semba, K., Watanabe, T. and Wada, H., En bloc immunohistochemistry reveals extensive distributions of histidine decarboxylase-immunoreactive neurons on the ventral surface of the rat hypothalamus, 137
Rey, C., see Benistant, C., 237
Ríos, C. and Tapia, R., Changes in lipid peroxidation induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and 1-methyl-4-phenylpyridinium in mouse brain homogenates, 321
Röhrenbeck, J., Wässle, H. and Heizmann, C.W., Immunocytochemical labelling of horizontal cells in mammalian retina using antibodies against calcium-binding proteins, 255
Ronco, P., see Pollard, H., 267

Sabel, B.A., DelMastro, R., Dunbar, G.L. and Stein, D.G., Reduction of anterograde degeneration in brain damaged rats by GM_1-gangliosides, 360
Santangelo, F., see Licata, F., 293
Sato, A., see Kurosawa, M., 193
Sato, Y., see Kurosawa, M., 193
Savoldi, F., see Fussi, F., 308
Scherman, D., Janssen, P.F.M. and Laduron, P.M., Axonal transport of the synaptic vesicle monoamine carrier identified by [^{3}H]dihydrotetrabenazine binding in rat sciatic nerve, 231
Schiebe, M., see Walter, Chr., 209
Schwartz, J.-C., see Pollard, H., 267
Seller, H., see Dembowsky, K., 31
Semba, K., see Reiner, P.B., 137
Shiells, R.A. and Falk, G., Joro spider venom: glutamate agonist and antagonist on the rod retina of the dogfish, 221
Silveira, R., see Dajas, F., 87
Smart, T.G., see Marrion, N.V., 55
Smith, M.E., see Adham, N., 283
Sonntag, R. and Fritzsch, B., The development of the amphibian trochlear nucleus. An HRP study, 143
Stein, D.G., see Sabel, B.A., 360
Stratford, T.R., see Wirtshafter, D., 261
Suzuki, H., see Kurosawa, M., 193
Suzuki, H., see Naoi, M., 215
Székely, G. and Matesz, C., Trigeminal motoneurons with disparate dentritic geometry innervate different muscle groups in the frog, 161

Takano, Y., see Nagashima, A., 103
Tapia, R., see Ríos, C., 321
Thomas, W.E., see Jordan, F.L., 249
Tokimasa, T., see Tsurusaki, M., 303
Trottier, S., see Esclapez, M., 131
Tsurusaki, M., Tokimasa, T. and Akasu, T., Postsynaptic inhibition of the frog neuromuscular transmission by prostaglandin E_1, 303

Vale, W., see Cintra, A., 25
Valentini, M., see Facchinetti, F., 249
Van den Buuse, M. and De Jong, W., Grooming behavior of spontaneously hypertensive rats, 71
Verroust, P., see Pollard, H., 267
Vlachová, V., see Bolotina, V., 298
Voronin, L.L., see Hess, G., 187
Vyskočil, F., see Bolotina, V., 298

Wada, H., see Reiner, P.B., 137
Walter, Chr. and Schiebe, M., FMRF-NH_2-like factor from neurohaemal organ modulates neuromuscular transmission in the locust, 209
Wässle, H., see Röhrenbeck, J., 255
Watanabe, T., see Reiner, P.B., 137
Widerlöv, E., Bissette, G. and Nemeroff, C.B., Centrally administered neurotensin inhibites pentobarbital metabolism in mice but not in rats, 311
Wikström, A.-C., see Cintra, A., 25
Wirtshafter, D., Stratford, T.R. and Asin, K.E., Evidence that serotonergic projections to the substantia nigra in the rat arise in the dorsal, but not the median, raphe nucleus, 261
Wong, T.M. and Lee, A.Y.S., Chronic morphine treatment reduces the incidence of ventricular arrhythmias in the isolated rat heart induced by dynorphine$_{1-13}$ or myocardial ischemia and reperfusion, 61
Wood, P.L., see Iyengar, S., 226

Yoshioka, H., Nishino, K., Miyake, T., Ohshio, G., Kimura, T. and Hamashima, Y., Immunohistochemical localization of a new thiamine diphosphate-binding protein in the rat nervous system, 10

Zhang, L. and Krnjević, K., Effects of intracellular injections of phorbol ester and protein kinase C on cat spinal motoneurons in vivo, 287